

WHERE IS THE MATTER IN THE MERGING CLUSTER ABELL 2218?

ÁRDÍS ELÍASDÓTTIR², MARCEAU LIMOUSIN², JOHAN RICHARD³, JENS HJORTH², JEAN-PAUL KNEIB⁴, PRIYA NATARAJAN^{5,6},
 KRISTIAN PEDERSEN², ERIC JULLO⁷, DANUTA PARAFICZ^{2,8}

Draft version February 2, 2008

ABSTRACT

We present a parametric strong lensing model of the cluster Abell 2218 based on HST ACS data. We constrain the lens model using 8 previously known multiply imaged systems, of which 7 have spectroscopically confirmed redshifts. In addition, we propose five candidate multiply imaged systems and estimate their redshifts using our mass model. The model parameters are optimized in the source plane by a bayesian Monte Carlo Markov Chain as implemented in the publicly available software Lenstool. We find $\text{rms}_s = 0''.12$ for the scatter of the sources in the source plane, which translates into $\text{rms}_i = 1''.49$ between the predicted and measured image positions in the image plane. We find that the projected mass distribution of Abell 2218 is bimodal, which is supported by an analysis of the light distribution. We also find evidence for two structures in velocity space, separated by $\sim 1000 \text{ km s}^{-1}$, corresponding to the two large scale dark matter clumps. We find that the lensing constraints can not be well reproduced using only dark matter halos associated with the cluster galaxies, but that the dark matter is required to be smoothly distributed in large scale halos. At $100''$ (291 kpc) the enclosed projected mass is $3.8 \times 10^{14} M_\odot$. At that radius, the large scale halos contribute $\sim 85\%$ of the mass, the brightest central galaxy (BCG) contributes $\sim 9\%$ while the remaining $\sim 6\%$ come from the other cluster galaxies. We find that the model is not very sensitive to the fainter (and therefore by assumption less massive) galaxy sized halos, unless they locally perturb a given multiply imaged system. Therefore, dark galaxy sized substructure can be reliably constrained only if it locally perturbs one of the systems. The massive BCG and galaxies which locally perturb a multiply imaged system are reliably detected in the mass reconstruction. In an appendix we give a self-contained description of the parametric profile we use, the dual pseudo isothermal elliptical mass distribution (dPIE). This profile is a two component pseudo isothermal mass distribution (PIEMD) with both a core radii and a scale radii.

Subject headings: dark matter — galaxies: clusters: individual (Abell 2218) — gravitational lensing

1. INTRODUCTION

Dark matter dominates over baryonic matter in the universe, but its nature is not known. The study of the inner parts of dark matter halos can give insight into the nature of the dark matter, as the steepness of the profile is correlated with the interaction between the dark matter itself and with the baryonic matter. According to Λ CDM simulations, the mass distribution of galaxy clusters should be dominated by their dark matter halos. Gravitational lensing, which is sensitive to the total matter distribution, visible or dark, is ideal for studying the mass distribution of clusters. Strong lensing features, consisting of multiply imaged and strongly distorted background

sources, provide constraints on the inner parts of the cluster, while weak lensing features, consisting of weakly distorted singly imaged background sources, provide constraints on the outer slope of the surface mass profile (see e.g., Smail & Dickinson 1995; Smail et al. 1995; Kneib et al. 1996; Smail et al. 1997; Abdelsalam et al. 1998; Natarajan et al. 2002b; Bradač et al. 2006; Gavazzi & Soucaill 2007; Limousin et al. 2007b).

Lensing can therefore provide unique information about the total mass distribution of clusters, from the inner to the outer parts. In addition, lensing can in principle be used to deduce various cosmological parameters (e.g. H_0 , Ω_Λ , Ω_m). This has already been extensively applied to lensing on galaxy scales (see e.g., Schechter et al. 1997; Koopmans et al. 2003) and to a smaller degree for lensing on cluster scales (see e.g., Soucaill et al. 2004; Meneghetti et al. 2005), where the accuracy of the mass map is a limiting factor. The accuracy of the mass map is strongly dependent on the number of multiply imaged systems used to constrain it. Therefore, to construct a robust model of the dark matter distribution, accurate enough for cosmography and for using the cluster as a gravitational telescope, it is important to include as many spectroscopically confirmed multiply imaged systems as possible (see e.g., Ellis et al. 2001).

Abell 2218 is one of the richest clusters in the Abell galaxy cluster catalog (Abell 1958; Abell et al. 1989) and has been successfully exploited as a gravitational lens. A parametric lens model has previously been

Electronic address: ardis@dark-cosmology.dk

² Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Juliane Maries Vej 30, DK-2100 Copenhagen Ø, Denmark; (ardis, marceau, danutas, kp, jens)@dark-cosmology.dk.

³ Department of Astronomy, California Institute of Technology, 105-24, Pasadena, CA 91125; 5(johan,kneib)@astro.caltech.edu

⁴ OAMP, Laboratoire d'Astrophysique de Marseille UMR 6110 Traverse du Siphon 13012 Marseille, France; Jean-Paul.Kneib@oamp.fr

⁵ Astronomy Department, Yale University, P.O. Box 208101, New Haven, CT 06520-8101, USA, priya@astro.yale.edu

⁶ Department of Physics, Yale University, P.O. Box 208101, New Haven, CT 06520-8101; USA

⁷ European Southern Observatory, Alonso de Cordova 3107, Vitacura, Chile; ejullo@eso.org

⁸ Nordic Optical Telescope (NOT), Apartado 474, 38700 Santa Cruz de La Palma, Canary Islands, Spain

constructed by Kneib et al. (1995, 1996) (using 1 and 2 spectroscopically confirmed systems respectively) and by Natarajan et al. (2002b, 2007) (using 4 and 5 spectroscopically confirmed systems) building on the model of Kneib et al. (1996) and including weak lensing constraints from HST WFPC2 data. A non-parametric model was constructed by Abdelsalam et al. (1998) using three spectroscopically confirmed multiply imaged systems. In all of these models, a bimodal mass distribution was required to explain the image configurations (i.e. the models include two large scale dark matter clumps), but the number of constraints were not sufficient to accurately constrain the second large scale dark matter clump. Abell 2218 has also been used as a gravitational telescope, with Ellis et al. (2001) discovering a source at $z = 5.6$ and Kneib et al. (2004a) discovering an even more distant source at $z \sim 6.7$, later confirmed by Egami et al. (2005) using Spitzer data. Soucail et al. (2004) estimated cosmological parameters based on a lensing model of Abell 2218 using 4 multiply imaged systems. The latest published lens model of Abell 2218 is by Smith et al. (2005), who incorporated four multiply imaged systems and weak lensing constraints, using the WFPC2 data. Although the number of constraints has increased from the initial models, all previous models have assumed that the location of the second dark matter clump coincided with the brightest galaxy in the South East, due to a lack of constraints in its vicinity.

The motivation for revisiting the modeling of Abell 2218 comes from the new ACS data which have not been used before for modeling this cluster and are superior in both resolution, sensitivity and field of view to the previous WFPC2 data set. These new high quality data allow us to identify several subcomponents in previously known multiple images, thus adding more constraints, and in one case, two more multiple images of a known system. In addition, we have measured a spectroscopic redshift for an arc around the second dark matter clump, which our model predicts to be singly imaged. We also have several new candidate multiply imaged systems, which we add as constraints and estimate their redshift with the model. In total we have 7 multiply imaged systems with measured spectroscopic redshift and 6 multiply imaged systems without spectroscopic data (of which 5 are new candidate systems). Finally, the lensing code Lenstool, has undergone significant improvements from previous models, with the incorporation of a Monte Carlo Markov Chain (MCMC), which enables us to not only find the best model in the lowest χ^2 sense, but the most likely model as measured by its Evidence (Jullo et al. 2007). The MCMC also allows for a reliable estimate of the uncertainties in the derived model parameters.

The paper is organized as follows: In Section 2, we give an overview of the data used in this paper. We compile a list of all currently known and new multiply imaged systems in Abell 2218 and discuss the reliability of the redshift estimate of each one in Section 3. The methodology of the strong lensing modeling is described in Section 4. In Section 5 we present the results of our lensing analysis, and compare them to previous models. In Section 6 we discuss degeneracies in the modeling. In Section 7 we address how reliable our model is, and discuss the smoothness of the dark matter distribution. In Section 8 we interpret the bimodality of our model, along




FIG. 1.— Color-magnitude diagrams, (V-Z) and (I-Z), used for selecting the cluster members (see Section 2.1). The red quadrilaterals define the red sequence and galaxies lying within are considered cluster members.

with X-ray measurements and an analysis of the distribution of cluster members in velocity space. We summarize our main conclusions in Section 9. Throughout the paper, we adopt a flat Λ -dominated Universe with $\Omega_\Lambda = 0.7$, $\Omega_m = 0.3$ and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$. Following Smith et al. (2005) we place the cluster at $z = 0.171$. At this redshift $1''$ corresponds to 2.91 kpc in the given cosmology.

2. DATA

We use data from several different sources for our lens modeling and analysis. The basis of our modeling is Advanced Camera for Surveys (ACS) data from the Hubble Space Telescope, which allows us to identify and accurately locate multiply imaged systems. Our cluster member catalog is also selected using the ACS data and the magnitude of each cluster member is given in the Ingrid K-band of the William Herschel Telescope. In addition, we have obtained a spectroscopic redshift of a system using the Keck telescope and archival Chandra X-ray Observatory data have been used to produce an X-ray map of Abell 2218.

2.1. The galaxy catalog

Cluster members were selected based on the ACS data using the characteristic cluster red-sequences (V-Z) and (I-Z) in two color-magnitude diagrams and selecting objects lying within the red-sequences of both diagrams (see Figure 1). The ACS data were used to select the galaxy members due to their better photometric accuracy compared to the K-band data. Nevertheless, the catalog is given in the K band since it is more representative of the galaxy population in elliptical galaxies. This selection gave 203 cluster galaxies down to $K = 19.6$. However, we rejected six galaxies from the catalog leaving 197 cluster members, of which four have measured redshifts showing them to be background galaxies.

2.2. X-ray data

The X-ray data were taken with the Advanced CCD Imaging Spectrometer (ACIS) on NASA's Chandra X-ray Observatory. The ACIS is an array of 10 CCD's, capable of simultaneously imaging and measuring the energy of the incoming X-rays. We combined three images




FIG. 2.— A color image of Abell 2218 based on ACS data (F775W, F625W and F475W filters in the red, green and blue channel respectively). The cluster galaxies are marked in yellow (modeled using scaling relations) or blue (individually fitted). The multiple images are labeled in green for spectroscopically confirmed systems and red for candidate systems. The arc for which we have obtained spectroscopic redshift, S8, is labelled in cyan. Also shown are the critical lines corresponding to $z = 0.702$ (cyan), $z = 2.515$ (red) and $z = 6.7$ (green).

taken at different epochs to construct our final X-ray map of Abell 2218. Two of the images were taken in October 1999 and one in August 2001, with 5960 s, 11560 s and 49240 s exposure times respectively. The data reduction was performed using standard pipelines from the Chandra Interactive Analysis of Observations (CIAO) software. The X-ray data will be used for comparison with the lensing data in Section 8.

3. MULTIPLY IMAGED SYSTEMS

Multiply imaged systems are the basis of our analysis, and are particularly useful as constraints if their redshifts are accurately determined. In this section we list the multiple images, with or without spectroscopically confirmed redshifts, used as constraints in our lens modeling. We propose several new candidate systems found in the ACS data and present a spectroscopic redshift measurement for an arc which lies in the region of the second dark matter clump. All systems are listed in Table 1 and shown on Figure 2.

3.1. Previously known systems

The first system, system S1, is at a redshift of $z = 0.702$ as measured by Le Borgne et al. (1992) (system 359 in their catalog). Thanks to the new ACS observations, it has seven identified images (previously it had five identified images (Kneib et al. 1996)), of which two pairs are located close to individual galaxies #1028 and #993 in the galaxy catalog) which affect the lensing signal.

System S2 is a star-forming galaxy at a redshift of $z = 2.515$ (Ebbels et al. 1996). It consists of a fold and a counter image (listed as images #384 and #468 in Ebbels et al. (1996)). We separate the fold image into two images and identify four components, giving us four sets of triple images from this system (referred to as S2.1, S2.2, S2.3 and S2.4).

System S3 is a faint star-forming doubly imaged system at a high redshift $z = 5.576$ discovered by Ellis et al. (2001).

System S4 is a triply imaged submillimeter source at $z = 2.515$ reported by Kneib et al. (2004b). In each image they identify three components labelled α, γ, β . We identify all three images in the ACS data. However, we only detect the α and β components. We include both as constraints on the model (referring to them as S4.1 and S4.2 respectively). Image S4.1 seems to consist of further three subcomponents. As we do not clearly distinguish them in all three images, we do not add them as separate constraints.

The fifth system, S5, is a triply imaged system, believed to be the triply imaged outskirts of an associated singly imaged galaxy. It is believed that the galaxy partially crosses the caustic, with the central part being singly imaged and the outskirts being triply imaged. Kneib et al. (2004a) measured a redshift of $z = 2.515$ for the galaxy (labelled as #273 in their notation) but an independent spectroscopic redshift has not been determined for the faint triply imaged component. Although

TABLE 1
LENSED SYSTEMS

System id	R.A (J2000)	Dec. (J2000)	$z^\dagger$	Comments
S1.a	248.96976	66.212219	0.702	(1)
S1.b	248.96646	66.208627	0.702	
S1.c	248.96478	66.209439	0.702	
S1.d	248.96218	66.207296	0.702	
S1.e	248.95876	66.206237	0.702	
S1.f	248.95790	66.206762	0.702	
S1.g	248.95500	66.205771	0.702	
S2.1.a	248.95540	66.218525	2.515	(2)
S2.1.b	248.95451	66.218469	2.515	
S2.1.c	248.93684	66.206345	2.515	
S2.2.a	248.95562	66.218536	2.515	
S2.2.b	248.95426	66.218458	2.515	
S2.2.c	248.93681	66.206424	2.515	
S2.3.a	248.95642	66.218428	2.515	
S2.3.b	248.95441	66.218347	2.515	(3)
S2.3.c	248.93723	66.206355	2.515	
S2.4.a	248.95405	66.218258	2.515	
S2.4.b	248.95715	66.218375	2.515	
S2.4.c	248.93728	66.206269	2.515	
S3.a	248.96646	66.214324	5.576	(4)
S3.b	248.96591	66.212701	5.576	
S4.1.a	248.97894	66.210224	2.515	
S4.1.b	248.97519	66.206897	2.515	
S4.1.c	248.96075	66.201462	2.515	
S4.2.a	248.97981	66.210279	2.515	
S4.2.b	248.97463	66.206179	2.515	
S4.2.c	248.96296	66.201575	2.515	(5)
S5.a	248.981043	66.205816	2.515	
S5.b	248.977680	66.203939	2.515	
S5.c	248.974235	66.202425	2.515	
S6.a	248.9855	66.200483	1.034	
S6.b	248.98371	66.199906	1.034	
S7.a	248.94254	66.216974	1.03	
S7.b	248.94244	66.216909	1.03	(6)
S7.c	248.95763	66.220132	1.03	
S8.a	248.98805	66.19633	2.74	
C1.a	248.97735	66.210845	~ 6.7	
C1.b	248.97585	66.208989	~ 6.7	
C1.c	248.95303	66.200703	~ 6.7	
C2.a	248.95398	66.203803	2.6 ± 0.1	
C2.b	248.96247	66.204748	2.6 ± 0.1	
C2.c	248.97692	66.216365	2.6 ± 0.1	
C3.a	248.93879	66.22238	2.8 ± 0.6	
C3.b	248.93749	66.221662	2.8 ± 0.6	
C4.a	248.97708	66.210108	2.2 ± 0.2	
C4.b	248.97542	66.208369	2.2 ± 0.2	
C5.a	248.97543	66.209244	2.3 ± 0.8	
C5.b	248.97556	66.209383	2.3 ± 0.8	
C6.a	248.96481	66.213039	2.6 ± 0.3	
C6.b	248.96505	66.214017	2.6 ± 0.3	

NOTE. — The multiply imaged systems used to constrain the model. The systems and their properties, along with references to the papers reporting their redshifts, are given in Section 3. Systems C2-C6 are new candidate systems reported in this paper. Systems with a spectroscopic redshift measurement are denoted by an S in their id, while those without spectroscopic redshifts are denoted by a C. Roman letters (a,b,c,...) denote the different images of each set of multiple images. If several components are identified in system, each component is sublabelled, e.g. 2.1 and 2.2 denote two different components of the same multiple imaged system. (1) Two pairs of images are caused by splitting by a local cluster galaxy. (2) Also known in the literature as #384 and #486. (3) The redshift is measured for an associated singly imaged galaxy (4) Also known in the literature as #289 (5) Also known as #444 and H6. The redshift estimate is based on one feature. (6) Redshift estimate based on photometric redshift, lensing and a tentative Ly α break.

† Lens model redshift predictions are reported in square brackets.




FIG. 3.— The new candidate system, C2, identified in the ACS images. The three panels show the three images of C2, marked in Figure 2. The colors are the same as in Figure 2. The system is characterized by a central 'yellow' spot, flanked by blue spots.




FIG. 4.— The new candidate arcs identified in the ACS images. *Left panel:* The candidate arc C3 (green circles). *Middle panel:* The candidate systems C4 (green circles) and C5 (yellow circle). The red circles are the locations of two of the images of system C1 (at $z = 6.7$), while the blue circle shows the location of images belonging to system S4 (components a and b). *Right panel:* The candidate system C6 (green circles). The high redshift system S3 ($z = 5.6$) is marked with red circles.

it is likely (and consistent with the models) that they belong to the same background source, it is not certain, making the redshift determination for S5 less certain.

The sixth system, S6, consists of an arc in two parts at $z = 1.034$ (Kneib et al. 1996; Swinbank et al. 2003, called 289 in their notation).

The seventh system, S7, first reported by Kneib et al. (1996), consists of a merging image and a counter image (also known as #444 and H6 respectively). Ebbels et al. (1998) measured a spectroscopic redshift at $z = 1.03$ for the merging image but this is listed as a tentative identification (as the only identified feature in the spectra was the Fe II doublet at $\lambda\lambda 2587, 2600$).

The final previously known multiply imaged system, C1, is a high redshift ($z \sim 7$) triply imaged system (Kneib et al. 2004a). Egami et al. (2005) further constrained the redshift of the system to lie in the range 6.6 – 6.8. For the purposes of the modeling we will assign $z = 6.7$ to this system, noting that changing the redshift within the range 6.6 – 6.8 does not affect the model.

3.2. New candidate systems

We propose 5 new candidate systems which we have identified in the new ACS data. We do not have spectroscopic redshift determination for these systems, but estimate their redshifts using the model prediction.

The first candidate is a triply imaged system, which we call C2. In Figure 3 we show color stamps of the three images, and the locations are given in Table 1. The morphology of the three images, suggests that it is of the same background source, with a 'yellow' spot in the middle, flanked by a fainter blue on the sides.

We also identify a new faint arc in the north-west part of the cluster, which we call C3 (see Figure 4). In the vicinity of system 6, we further identify two arclike images, which we label C4 and C5. The first, C4, is a very faint blue double arc, while in the second, C5, two bright spots of a merging arc can be seen. Finally, a pair of blue extended images, C6, is found in the vicinity of system 3. These candidate systems are shown in a color




FIG. 5.— The alignment of the slit for the spectra of S8. The box shows the area used to extract the spectrum (see Figure 6).




FIG. 6.— The spectrum of system S8. The plot shows lines for the arc at $z = 2.74$ (blue) and the nearby galaxy (#617) at $z = 0.177$ (red). The spectrum shows Ly α absorption and emission and a number of absorption lines in the UV which are marked in the figure. At this redshift, $z = 2.74$, the model predicts S8 to be singly imaged.

image in Figure 4 and are listed in Table 1 along with their redshift estimates (see Section 5.4 for details). The positions of these images are used to constrain the final model, but their redshift is kept free in the modeling.

3.3. Spectroscopic redshift for an arc

We have used the Low Resolution Imager and Spectrograph (LRIS, Oke et al. (1995)) at Keck to measure the spectroscopic redshift of an arc, S8, to the south-east of the second dark matter clump (see Figure 2). Two exposures of 1800 seconds were obtained on June 29th 2007 with a $175'' \times 1''$ long slit placed along the brighter components of this arc (Figure 5). A 6001 mm^{-1} grism blazed at 4000 Å and a 400 mm^{-1} grating blazed at 8500 Å were used in the blue and red channels of the instrument, both lightpaths being separated by a dichroic at 5600 Å . The corresponding dispersions are $0.6/1.85 \text{ Å}$ and resolutions are $4.0/6.5 \text{ Å}$ in the blue/red channel, respectively.

The resulting spectrum (see Figure 6) is dominated by the light coming from the very bright neighbor cluster member at $z = 0.177$, yet shows Lyman- α in emission and UV absorption features of SiII, CII and CIV from the arc, giving a redshift $z = 2.74$. These features are not compatible with any line at the redshift of the lens galaxy, giving a redshift class of 2 for this spectrum, with 75% probability of being correct, following the classification of Le Fevre et al. (1995). Our model is not consistent with this object being multiply imaged,

and when including it as a constraint as multiply imaged (using spots in the arclike structure as independent images), the model predicts additional counter images which are not seen. We therefore conclude that although this arc is lensed, it is not multiply imaged, and include it as a singly imaged system in the model constraints.

4. MODELING

Following e.g., Kneib et al. (1996); Smith et al. (2005); Limousin et al. (2007b) we do a parametric mass reconstruction of Abell 2218. The multiply imaged systems form the basis of our analysis, with each n -tuply imaged system giving $2(n-1)$ constraints if the redshift is known. As the number of constraints needs to be greater than the number of free parameters in our fit, we are limited by the known multiply imaged systems. The MCMC sampling and optimization is done using the Lenstool software (Kneib 1993; Jullo et al. 2007). The optimization is performed in the source plane, as it is faster and has been found to be equivalent to optimizing in the image plane (Halkola et al. 2006; Jullo et al. 2007). The new version of Lenstool (Jullo et al. 2007), gives a distribution of values for each of the parameters, thus making it possible to estimate the uncertainty of the parameters. In addition, it returns the Evidence of a model, which is a measure of how likely a model is, penalizing unnecessarily complicated models. Thus, a model with a lower χ^2 but more free parameters, may have a lower Evidence, suggesting that we should choose the simpler model. For each image we find its rms (root-mean-square) value for its position in the source plane, rms_s , and image plane, rms_i , given by

$$\text{rms}_s = \sqrt{\frac{1}{n} \sum_{j=1}^n \left(X_s^j - \langle X_s^j \rangle \right)^2} \quad (1)$$

$$\text{rms}_i = \sqrt{\frac{1}{n} \sum_{j=1}^n \left(X_{\text{obs}}^j - X^j \right)^2} \quad (2)$$

where n is the number of images for the system, X_s is the position in the source plane, X the position in the image plane and X_{obs} the observed position in the image plane. The overall rms is defined by summing and averaging over all the images for all the systems. A detailed overview of the Lenstool software, including definitions of χ^2 and the Evidence, can be found in Jullo et al. (2007).

4.1. Model components

We refer to the individual components of the model as 'clumps', where each clump is denoted by its position, ellipticity, position angle and the parameters of the profile used to describe it. The parametric profile we use is the dual Pseudo Isothermal Elliptical mass distribution (dPIE, derived from Kassiola & Kovner (1993b)) and its form and main properties are given in Appendix A. The dPIE profile is defined in Lenstool by three parameters, the core radius a , the scale radius s and a fiducial velocity dispersion σ_{dPIE} . For $a < r < s$, the profile behaves as $\rho \sim r^{-2}$, while it falls like r^{-4} in the outer regions. For a vanishing core radius the scale radius corresponds to the radius containing half the 3D mass.

The clumps are called 'large scale clumps' if their mass within the outermost multiply imaged constraint

is greater than 20% of the total mass. Smaller clumps are referred to as 'galaxy scale clumps', and are in general associated with the cluster members. Large scale dark matter clumps are optimized independently. The redshift is fixed at the location of the cluster, but the central position (R.A., Dec.), the ellipticity and the position angle (P.A.) are allowed to vary.

We associate a galaxy scaled clump with each of the cluster galaxies, fixing the central location, ellipticity and position angle of the mass distribution to that of the light distribution. A few of the cluster galaxies are fitted individually (optimizing their a , s and σ_{dPIE} , see Table 2 for an overview), but due to a lack of constraints, most of the galaxies are optimized in a combined way. The parameters (a , s , σ_{dPIE}) are optimized together using the following scaling relations for the luminosity L :

$$a = a^* \left(\frac{L}{L^*} \right)^{1/2} \quad (3)$$

$$s = s^* \left(\frac{L}{L^*} \right)^{1/2} \quad (4)$$

$$\sigma_{\text{dPIE}} = \sigma_{\text{dPIE}}^* \left(\frac{L}{L^*} \right)^{1/4}. \quad (5)$$

For a discussion of these scaling relations we refer to Limousin et al. (2007b) and Halkola et al. (2007). For a given L^* luminosity, we fix $a^* = 0.25$ kpc, while σ_{dPIE}^* and s^* are allowed to vary. We note that fixing the core radius to be small, makes the profile approximately equivalent to the profile used by Brainerd et al. (1996) to describe galaxies (see also Appendix A). Following de Propris et al. (1999) we take the apparent magnitude of an L^* in the K-band to be $K^* = 15$ at $z = 0.171$ (redshift of Abell 2218).

5. THE STRONG LENSING MASS DISTRIBUTION

In this section we present our strong lensing model (optimized in the source plane), discuss its implications and compare it to previous results. All reported error bars correspond to 68% confidence levels. For our best model we find $\text{rms}_s = 0''.12$, which gives $\text{rms}_i = 1''.49$ (see Table reftab:chires).⁸

5.1. A bimodal mass distribution

We show the mass density model in Figure 7 and critical lines predicted by the model at $z = 0.702, 2.515, 6.7$ in Figure 2. The total projected mass as a function of radius, centered on the BCG, is shown in Figure 8. We find that the mass distribution is strongly preferred to be bimodal, even for a simple model where all the galaxies are modelled based on the scaling relations, but the dark matter clumps are optimized independently using only systems 1 and 2 as constraints (28 constraints in all). The second dark matter clump is located to the south east of the central clump for this simple model, near galaxy #617, which is in agreement with previous

models of Abell 2218. The bimodal mass distribution is also strongly preferred when we add more constraints. We will refer to the two large scale dark matter clumps as DM1 (for the one associated with the BCG) and DM2 (for the one associated with galaxy #617). We also constructed a three clump model, but its bayesian Evidence was worse, leading us to reject it as our best model.

In previous models, the location of the second clump has been fixed at the center of the brightest galaxy in the south east corner (#617), but we find that when the location is allowed to vary it is offset from this galaxy by $\sim 35''$. We also find that DM2 has high ellipticity and is comparable in mass to DM1, although significantly less massive than the DM1 and BCG halos combined (see Figure 8). We note that the light distribution is similar to the derived matter distribution (see Figure 7), supporting the finding that there is a significant matter component in the vicinity of DM2 (see also section 8).

5.2. Dark matter halos of galaxies

The potential contribution of the individual dark matter halos associated with cluster galaxies was first proposed by Natarajan & Kneib (1997). Typically it is found that contribution of dark matter halos associated with the bright, early-type cluster galaxies in the inner regions are required to explain the positions and geometries of multiply lensed images in the strong lensing regime (Meneghetti et al. 2007).

We fit three galaxies individually (for the parameters of these galaxies see Table 2), two are the brightest galaxies near the centers of the large scale dark matter clumps (#1193 - the BCG - and #617), while the third is an elliptical galaxy important to the lensing of system S1. The halo associated with the BCG is very massive, and in particular in the inner $10''$ it dominates the mass distribution (see Figure 8). Even at the outermost Einstein radii of a multiply imaged system ($80''$) it still contributes around 10% of the mass of the cluster.

System S1 has seven images, and two of the pairs are strongly affected by nearby galaxies (an elliptical #1028 and a spiral #993). Keeping the parameters of the spiral free did not significantly affect the model, while adding the elliptical did. Although both galaxies would at first glance appear equally important to be fitted individually due to their strong effects on system S1, Figure 9 provides an important insight to why only #1028 needs to be included individually. This is because the scaling relations (see § 4.1) assume that the galaxies follow a Faber-Jackson relation (Equation 5) without any scatter. As can be seen from Figure 9 and discussed in Ziegler et al. (2001) the galaxies in Abell 2218 show a significant scatter and therefore the above scaling relations can give very unrealistic values for individual galaxies which lie far from the mean behavior. The elliptical #1028 is one such galaxy, with a measured central velocity dispersion which is greater than the central velocity dispersion of the BCG as measured by Jørgensen et al. (1999), while #993, although a spiral galaxy, is more consistent with the mean.

For the other cluster galaxies, they are included in the model via the scaling relations given in Section 4.1. For L^* corresponding to $K = 15$ at $z = 0.171$ we find $\sigma_{\text{dPIE}}^* = 185_{-11}^{+10}$ km s⁻¹ and $s^* = 2''.9_{-0.3}^{+0.5}$, but we note that there is some degeneracy between the two values

⁸ A parameter file containing all the following information, and which can be used with the Lenstool software package, along with a FITS file of the mass map generated from the best-fit model are available at <http://archive.dark-cosmology.dk/>. These can be used to find relevant critical lines for using Abell 2218 as a gravitational telescope and to predict/confirm candidate lensed systems.

TABLE 2
MODEL PARAMETERS

Clump	Δ R.A. "	Δ Dec. "	$\hat{\epsilon}$	$\theta_{\hat{\epsilon}}$	σ_{dPIE} (km/s)	a (")	s (")
DM1	$3.9^{+0.3}_{-0.4}$	$23.1^{+1.7}_{-0.7}$	$0.033^{+0.03}_{-0.001\dagger}$	56^{+6}_{-6}	806^{+10}_{-10}	26^{+1}_{-1}	274^{+110}_{-4}
DM2	$-20.4^{+0.1}_{-0.6}$	$-23.1^{+0.6}_{-0.6}$	$0.39^{+0.01\dagger}_{-0.01}$	8^{+1}_{-2}	1029^{+13}_{-49}	44^{+2}_{-3}	462^{+85}_{-60}
1193 (BCG)	-0.5	0.07	0.46	52.4	514^{+7}_{-11}	$3.14^{+0.04}_{-0.45}$	56^{+12}_{-8}
617	-46.1	-49.1	0.20	59.4	288^{+7}_{-8}	$2.5^{+0.3}_{-0.5}$	87^{+10}_{-9}
1028	-16.0	-10.3	0.18	80.4	396^{+13}_{-128}	$0.3^{+0.2}_{-0.1}$	$0.6^{+0.5}_{-0.1}$
					σ_{dPIE}^* (km/s)	a^* (")	s^* (")
L* galaxy	...	...	...	...	185^{+10}_{-11}	0.09	$2.9^{+0.5}_{-0.3}$

NOTE. — Values quoted without error bars were kept fixed in the optimization. The error bars correspond to 68% confidence levels. The location and the ellipticity of the matter clumps associated with the cluster galaxies were kept fixed according to the light distribution. The center is defined at (R.A., Dec.)=(248.9546, 66.2122) in J2000 coordinates.

$\dagger$ The posterior probability distribution pushes towards the highest values of our input prior. Therefore these error bars do not represent the full uncertainty in the value, but the uncertainty given a prior of $\hat{\epsilon} < 0.4$




FIG. 7.— The mass density map and its contours (black). The maps are $300'' \times 300''$ and centered on the BCG with North being up and East being left. The red contours show the light distribution (left panel) and the X-ray distribution (right panel). The overall shape of the light-contours and the mass map contours are similar, with the light being slightly more 'pear'-shaped, as it broadens in the SE direction. The overall shape of the light-contours and the mass map contours also agree, although the X-rays become more spherical for in the outer regions. See section 8.

(see discussion in section 6).

5.3. Comparison with measured velocity dispersions

We use the velocity dispersion measurements, σ_0 , of Ziegler et al. (2001) and Jørgensen et al. (1999) for Abell 2218 to compare with the results from the lens model. Ziegler et al. (2001) have a total of 48 galaxies in their sample, of which nearly half fall within the ACS image and are included in our galaxy catalogue, while the Jørgensen et al. (1999) data contains the BCG and seven other cluster members in our galaxy catalogue. Using the velocity dispersion measurements for Abell 2218 cluster members from Ziegler et al. (2001) we plot the Faber-Jackson relationship for the K-band data in Figure 9. We then calculate the mean and the standard deviation of galaxies with K-band magnitudes in the range from 14.8 to 15.2, excluding the individually fitted galaxy (galaxy #1028 in our catalog, or #1662 in the notation used by Ziegler et al. (2001)). This gives

$\sigma_{0,\text{ziegler}}^* \approx 195 \pm 35 \text{ km s}^{-1}$ for a typical $K=15$ galaxy.

In Figure 10 we plot σ_0 from Ziegler et al. (2001) and Jørgensen et al. (1999) vs. the fiducial velocity dispersion, σ_{dPIE} for the BCG, the two individually fitted galaxies and L^* . Although the values from direct velocity dispersion measurements can not be directly related to the values obtained from the dPIE profile (as the measured values are calculated based on an isothermal profile and are not aperture corrected) we find that they are consistent with σ_{dPIE} being related to the measured velocity dispersion by $\sigma_0 \approx 0.85\sigma_{\text{dPIE}}$ as found in Section A.7.

5.4. Redshift estimates of the new candidate systems

We estimated the redshifts of the new candidate systems using the model predictions. The three component candidate system C2 is found to have $z = 2.6 \pm 0.1$. For the merging arcs we find $z = 2.8 \pm 0.6$ for C3, $z = 2.2 \pm 0.2$ for C4, $z = 2.6 \pm 0.3$ for C6 while C5 is poorly constrained




FIG. 8.— Total projected mass as a function of aperture radius (centered on the BCG) for different model components. The two large scale clumps, DM1 and DM2, contribute a similar amount to the mass, but the halo associated with the BCG dominates in the inner regions, and the combined halo of DM1 and the BCG dominates the mass in the region where most of the constraints lie (within $80''$). The galaxies (excluding the BCG) only contribute a small amount to the overall mass, of the order of 5 – 6%.




FIG. 9.— The Faber-Jackson relation for the K-band using σ_0 measurements from Ziegler et al. (2001). Also plotted is the central velocity dispersion of the BCG (#1193) as obtained by Jørgensen et al. (1999). The individually marked galaxies are discussed in section 4.1 and 5.2.




FIG. 10.— The velocity dispersion of the dPIE, σ_{dPIE} vs. σ_0 measurements from Ziegler et al. (2001) and Jørgensen et al. (1999). The line is not a fit, but shows the relationship $\sigma_0 = 0.85\sigma_{\text{dPIE}}$ found in Section A.7.

TABLE 3
GOODNESS OF THE FIT

System	χ^2	rms_s ($''$)	rms_i ($''$)
S1	21.7	0.171	0.60
S2.1	10.0	0.090	0.60
S2.2	11.8	0.098	0.82
S2.3	37.9	0.167	1.49
S2.4	55.8	0.197	2.17
S3	4.9	0.043	0.57
S4.1	9.8	0.089	0.53
S4.2	2.1	0.042	0.43
S5	15.6	0.060	4.61
S6	0.3	0.016	0.03
S7	8.4	0.052	1.45
S8	0.0	0.000	0.00
C1	13.4	0.102	1.26
C2	20.5	0.178	0.63
C3	17.5	0.122	1.64
C4	4.7	0.052	0.70
C5	0.2	0.009	0.27
C6	0.5	0.011	0.14
	$\chi^2/\text{d.o.f.}$	rms_s	rms_i
Full model	235.1/34	0.12	1.49

NOTE. — The χ^2 and rms in the source plane (where the model is optimised) and the image plane. The zero values found for S8 mean that the system is consistent with being singly imaged.




FIG. 11.— A plot of the source plane at $z = 2.515$ showing systems S2, S4 and S5 and the caustic lines. Their maximum separation is around 130 kpc, consistent with them belonging to the same group of galaxies. Also shown are the candidate systems C2, C3, C4, C5 and C6 when their redshift is assumed to be $z = 2.515$. Under that assumption, their locations are consistent with them also belonging to this group of galaxies.

with $z = 2.3 \pm 0.8$. The estimated redshifts are reported in Table 1. These redshifts are consistent with a preliminary photometric redshift analysis done using the Hyperz photometric redshift code (Bolzonella et al. 2000).

5.5. A strongly lensed galaxy group at $z=2.5$

Three multiply imaged systems, S2, S4 and S5, all have the same redshift of $z = 2.515$. To check whether these three systems belong to a background galaxy group, we lens them back to their source plane (see Figure 11).

We find that S4 is in the middle with S5 and S2 at a separation of $5''.4$ and $10''.3$ respectively, with the maximum separation being $15''.7$, corresponding to 127 kpc in the source plane ($1'' = 8.06$ kpc at $z = 2.515$), suggesting that the three systems belong to the same background

group of galaxies. This is in agreement with the findings of Kneib et al. (2004b) who found the maximum separation of the systems in the source plane to be 130 kpc. It may be of interest to do a dedicated search for more systems at $z = 2.515$, either multiply imaged or singly imaged, to further study this high redshift group of galaxies, and we note that all the candidate systems have a predicted z consistent with $z = 2.515$. If we assume that these candidate systems have $z = 2.515$, their location in the source plane is consistent with them belonging to this same group (see Figure 11).

5.6. Comparison with previous results and weak lensing

Although the overall results of our model are in agreement with previous models of Abell 2218, they have found the second clump to be less massive (see e.g., Kneib et al. 1996; Abdelsalam et al. 1998; Smith et al. 2005; Natarajan et al. 2007). There are several possible reasons for the discrepancy, with the first one being the new constraints we use in this model. We have therefore redone the modeling using only the previously known spectroscopically confirmed systems, but we still find that DM2 is massive and with a large core. It is also possible, that the previous models constructed using the older version of the lenstool package which did not involve MCMC sampling, have been caught in local minima. Indeed, we do find when forcing DM2 to be smaller, a comparable χ^2 but with the posterior probability distribution of the core radius a pushing toward the upper limit of the input range. However, this can not explain the discrepancy found for the non-parametric model of Abdelsalam et al. (1998).

A third possible explanation is that the models of Abdelsalam et al. (1998); Smith et al. (2005); Natarajan et al. (2007) all incorporated weak lensing to ‘anchor’ the outer part of the mass distribution. Using our model to predict the weak lensing shear profile at large radii, we find that our profile overestimates the signal compared to the measured signal of Bardeau et al. (2007) from ground based observations with the Canada-France-Hawaii Telescope (see Figure 12). The central part of the Bardeau et al. (2007) profile is flat, characteristic of contamination of the background galaxy catalog by foreground cluster members (see Limousin et al. (2007b) for discussion on contamination). Therefore, we do not expect to have an agreement between the weak lensing result and the strong lensing result. At around $\sim 300 - 400''$ the contamination should be negligible, and there we find that the agreement is better although our model still overpredicts the signal (although they are consistent within 2σ). We stress however that at this radius, we are extrapolating a strong lensing model based on constraints within $100''$, and the prediction becomes more uncertain the further we go out.

6. DEGENERACIES

In Section 5 we presented the strong lensing model of Abell 2218. In this Section we study the degeneracies of our parametric strong lensing modeling, both those inherent to the parametric profile and the model components. Jullo et al. (2007) have also discussed the various degeneracies of the dPIE profile in lensing, addressing how different image configurations can break some degeneracies.




FIG. 12.— The weak lensing signal predicted by our model (solid line) compared to the weak lensing found by Bardeau et al. (2007) (dotted line - 1σ error bars). The Bardeau et al. (2007) shows a very flat inner profile, characteristic of contamination of the background galaxy catalog by cluster members. Therefore, we do not expect an agreement in the inner regions. In the outer regions, where the contamination should be negligible, we find that our model overpredicts the weak lensing signal, but we note that the prediction is an extrapolation of a strong lensing model with constraints within $100''$.




FIG. 13.— The degeneracy between the scale radius, s , and σ_{dPIE} for #1028 (left) and the scaled galaxies (right). These arise from equation A10 which gives the aperture mass, showing that to keep the enclosed mass constant, an increase in the scale radius requires a lower value of σ_{dPIE} . The contours correspond to 1σ , 2σ and 3σ confidence levels.

Lensing most strongly constrains the projected mass, and therefore we expect to see degeneracies arising from Equation A10, although ellipticity may complicate that picture further. In agreement with Jullo et al. (2007) we find for the large scale halos, that the scale radius, s , is poorly constrained (as it lies beyond the outermost multiply imaged system). This is also the case for the BCG and the #617 which have large scale radii. For the smaller halos, i.e., the scaled galaxies and #1029, the scale radius, s , is small enough to affect the projected mass, and we find that lower s requires higher σ_{dPIE} to keep the mass constant (see Figure 13). However, the favored region for s is always small, consistent with the tidal stripping of cluster galaxies proposed by Natarajan et al. (1998, 2002a,b); Limousin et al. (2007a,c).

Also, in agreement with Jullo et al. (2007), and as discussed by e.g. Kochanek (1996) for more general cored profiles, a larger core radius, a , requires higher σ_{dPIE} to keep the mass constant (see Figure 14). As our model has




FIG. 14.— The degeneracy between the core radius, a , and σ_{dPIE} for DM1, DM2, BCG, #1028, #617 from left to right. These again arise from Equation A10 requiring the aperture mass to remain constant. They show that to keep the mass constant, the σ_{dPIE} must increase when the core radius a is increased. This is consistent with the findings of Jullo et al. (2007) for the dPIE profile and the findings of Kochanek (1996) for more general profiles with core. The contours correspond to 1σ , 2σ and 3σ confidence levels.




FIG. 15.— The 2D posterior distribution of the parameters of DM2 vs. DM1 and BCG. There is a clear degeneracy between the parameters of the two large scale components and the BCG. Therefore, although the model prefers the inclusion of all the components, the values of their parameters are not fully independent. The contours correspond to 1σ , 2σ and 3σ confidence levels.

both a large core radius a and σ_{dPIE} for DM2 compared to previous models of A2218, we have explored whether this degeneracy can reduce both values. However, when forcing both a and σ_{dPIE} to be smaller, the posterior distribution of a always pushes to the upper limit of the input range. Therefore, we conclude that this degeneracy is not the explanation for the high a and σ_{dPIE} we find for DM2. This flat profile is further supported by the 'blind tests' we perform in Section 7.3.

In addition to the degeneracies associated with the profile itself, there may be degeneracies associated with the components we include in the model, i.e., are all the components we include necessary and are they independent of each other? As mentioned in section 5.1, the model is strongly preferred to be bimodal. There is however degeneracy between the parameters of DM1 (and the BCG) and DM2 which is visualized in Figure 15. As for the BCG, we find that the Evidence is marginally higher

FIG. 16.— The 2D posterior distribution of the scaled galaxy parameters (σ^* , s^*) with respect to the core radii of the large scale dark matter clumps DM1 and DM2. Although there is a degeneracy between the parameters, the scale radius s^* is well constrained at low values. The contours correspond to 1σ , 2σ and 3σ confidence levels.

when both the DM1 and BCG are included in the model, suggesting that the data are sufficient to model both separately. Therefore we conclude, that although the model prefers the inclusion of all three components, the values for their parameters are not fully independent. Finally, we look at the parameters for the scaled galaxies with respect to DM1 and DM2 (see Figure 16). Although there is degeneracy the preferred value, in particular for the scale radius s^* , is well constrained at low values ($\lesssim 4''$), consistent with the tidal stripping scenario.

7. RELIABILITY OF THE MASS MAP

We have presented the mass map inferred from our strong lensing analysis in Section 5 (Figure 7). In short, we find evidence for a bimodal mass distribution described by two large scale smooth dark matter clumps, on top of which we add some extra mass associated with the cluster galaxies. Moreover, we associate a significant mass concentration with the location of the BCG galaxy. These two conclusions are believed to be robust, and the aim of this section is to perform a series of tests in order to check our main findings regarding how the dark matter is distributed in Abell 2218.

7.1. Do we need the smoothly distributed dark matter component?

We wish to test whether the smoothly distributed component is necessary to reproduce a good fit, or whether




FIG. 17.— *Left panel:* The mass map from Section 7.1 using only halos associated with the galaxies. *Right panel:* The mass map from Section 5 consisting of large scale dark matter halos and galaxy halos. The mass map including the large scale dark matter halos, which is much smoother, provides a significantly better fit to the data than the clumpy galaxy-only model ($\text{rms}_s = 0''.12$ vs. $\text{rms}_s = 1''.62$). The maps are $300'' \times 300''$ and are centered on the BCG with North being up and East being left.




FIG. 18.— The total mass as function of radius (centered on the BCG) for the two models. Although the total mass within any given radii is similar, the pure-galaxy-mass map gives a significantly worse fit to the data ($\text{rms}_s = 0.12$ vs $\text{rms}_s = 1.62$), confirming the need for a large scale dark matter halo to accurately fit the data (see Section 7.1).

mass associated with the galaxies alone can provide an equally good fit. To this end, we construct a model where the mass is only in halos associated with the galaxies, without the inclusion of any smoothly distributed dark matter component. The galaxies are included in a scaled manner, allowing the σ_{dPIE}^* and s^* to move to higher values to increase their mass. As a result of the optimization, we find a very poor fit ($\text{rms}_s = 1''.62$ instead of $0''.12$ for the model from Section 5). As the model from Section 5 individually fits three of the galaxies, we need to check whether this difference arises from these extra free parameters. We therefore redo the analysis using only DM1 and DM2, and scaling all the cluster galaxies. The resulting fit is worse than the original ($\text{rms}_s = 0''.22$) but still significantly better than the fit without any smooth component.

The resulting mass map is shown in Figure 17 alongside our model from Section 5. The mass map without a smooth component is very 'clumpy', whereas the latter is smooth. As expected, the enclosed mass derived from

each model is comparable for any radius where we have observational constraints (see Figure 18). Therefore, the poorness of the 'clumpy' fit is not due to the fact that it is not massive enough to reproduce the lensing constraints. Moreover, it is worth noting that the 'clumpy' model is not very satisfactory in the sense that it describes the cluster galaxies as being very massive (around a few times $10^{12} M_\odot$ on average), which is not compatible with independent galaxy-galaxy lensing probes of cluster galaxy masses (see e.g., Mandelbaum et al. 2005). We therefore interpret the difference in the goodness of fit as evidence for the dark matter being distributed smoothly in the cluster, with only small perturbations from the cluster galaxies. This is further supported by the X-ray emission (see Section 8).

7.2. Sensitivity to the galaxy scale perturbations

We have already seen that individual galaxies are important to the overall lens model if they are close to a multiply imaged system (see also Section 7.3), but in general the cluster galaxies only add small perturbations to the overall mass distribution of the cluster, with ~ 5 -6% of the total mass being associated with the galaxy sized halos (excluding the BCG, see Figure 8). To check whether those small scale perturbations are important, we try using a subset of the galaxy catalog to see if this affects the results. Our original catalog contains 197 cluster members down to $K = 19.6$ (including the individually fitted galaxies), but we also create catalogs using a cut off magnitude of $K = 19, 18, 17, 16$ with 145, 110, 62, 35 members respectively. We find that the overall quality of the fit is only weakly affected (see Table 4), suggesting that for the purposes of lensing studies, it is sufficient to include the brightest galaxy members in the modeling in addition to those which clearly locally perturb given multiply imaged systems. This point is important in order to save computing time, as increased number of clumps, even if modelled by scaling, significantly increases the required CPU (Central Processing

TABLE 4
GOODNESS OF FIT FOR DIFFERENT CUT OFF MAGNITUDES

Cut off magnitude in K	# of galaxy members	$\log E$	rms_s (")	rms_i (")
19.6	197	-205	0.12	1.49
19.0	145	-192	0.12	1.05
18.0	110	-238	0.14	1.77
17.0	62	-284	0.17	1.87
16.0	35	-216	0.12	1.18

NOTE. — The χ^2 and Evidence (E) for different cutoff magnitude in the galaxy catalogue. All the models are optimised in the same way, the only difference being the number of galaxy members which are optimised using the scaling relations in section 4.1.

Unit) time.

7.3. Blind tests - can we localize galaxy scale substructure?

Normally galaxy sized perturbers are added to the model by placing a dark matter clump at the location of a known cluster member. This method is by construction not able to detect dark substructure directly. To test how sensitive we are to these substructures, luminous or dark, we perform 'blind tests', where we model the cluster with (A) five and (B) ten freely placed galaxy sized dark matter clumps, in addition to the two large scale dark matter clumps DM1 and DM2.

In order to limit the number of free parameters, we fix both $a = 1$ kpc and $s = 40$ kpc for the galaxy sized dark matter clumps, but we optimize all the parameters of the large scale clumps as before. We allow the positions of the galaxy scale clumps to vary from $-100''$ to $100''$, corresponding roughly to the ACS field of view. The velocity dispersion, σ_{dPIE} , of each clump is allowed to vary from 0-500 km s $^{-1}$.

The results are shown in Figure 19, showing a density plot of the position of the clumps for the 1000 best realizations (in the lowest χ^2 sense), for both configurations (A) and (B). The locations of the clumps corresponding to the best realization are marked by stars. We find that both configurations give a good fit to the data ($\text{rms}_s = 0.17$ and $\text{rms}_s = 0.19$ for (A) and (B) respectively). Configurations (A) and (B) are consistent with each other: setup (A) finds five well defined clumps. Setup (B) finds the same well defined five clumps, whereas it tries to marginalize the five extra clumps that seem not to be needed in the optimization.

For both configurations, the model places two large halos where the BCG is, clearly demonstrating the need for significant mass at the location of the BCG galaxy. Another common feature, is the lack of a galaxy sized clump in the vicinity of DM2, suggesting a very flat profile is preferred there. Both configurations place clumps in the positions of galaxies #1028 and #993, which are responsible for the local splitting of S1. This shows that lensing analysis can reliably detect galaxy sized dark substructure if it causes the local splitting of images, although the lensing map is not very sensitive to galaxy sized substructure in general (see also Section 7.2).

8. BIMODALITY OF THE MASS DISTRIBUTION: EVIDENCE OF A MERGER

Our strong lensing analysis shows that Abell 2218 has a bimodal mass distribution: even if the dominant mass

component is the BCG and DM1 halos, the second large scale dark matter clump DM2 is also significant, contributing around 20% of the total mass within $100''$. This second smooth mass component is associated with the bright cluster galaxy in the south-east that we call #617. To further interpret this bimodality, we look at two different available probes of the cluster: the X-ray map and the velocity of cluster members.

The X-ray map The X-ray flux map in the central parts of the cluster shows a complex morphology with no outstanding central peak. The offset between the X-ray peak at (R.A., Dec.)=(248.967, 66.210) and the BCG is significant, $\sim 20''$, with the X-ray peak located in the direction of DM2, but there is no evident peak in the X-ray emission in the vicinity of DM2. The X-ray flux map within about 1 arcmin of the peak of the X-ray emission is clearly elongated in the SE-NW direction, but becomes more spherical with increasing distance from the cluster center (see Figure 7). Comparing the contours for the X-rays and the mass map we see that the elongation of the two are very similar, although the X-rays become more spherical at large radii. To get more quantitative values for comparison of the two maps, we fitted a 2-dimensional β model (Cavaliere & Fusco-Femiano 1978) to both maps.

For the X-ray map, the best fit beta-model to the inner 2 arcmin by 2 arcmin (centered on the X-ray centroid) has an eccentricity ($\epsilon^\beta \equiv \sqrt{1 - (B/A)^2}$) of $\epsilon_X^\beta(\text{inner}) = 0.27 \pm 0.12$ and a position angle of $\theta_X(\text{inner}) = 39^\circ \pm 16^\circ$ (measured anti-clockwise from West). Fitting to the 4 arcmin by 4 arcmin X-ray map (centered on the X-ray centroid) results in eccentricity of $\epsilon_X^\beta(\text{outer}) = 0.18 \pm 0.08$ and a position angle of $\theta_X(\text{outer}) = 17^\circ \pm 14^\circ$. A similar analysis for the mass map in a 5 arcmin by 5 arcmin (centered on the BCG) gives an eccentricity of the overall mass map as $\epsilon_{\text{mass}}^\beta \approx 0.25$ with $\theta_{\text{mass}} \approx 39^\circ$, consistent with that found for the X-ray map, in particular in the inner regions.

Distribution in velocity space If a merger has taken place in Abell 2218, then this should be imprinted in the velocity distribution of the cluster members: we should be able to identify two structures in velocity space; one associated with the BCG galaxy, and one associated with galaxy #617.




FIG. 19.— Density plots for the positions of the galaxy sized clumps for the models discussed in section 7.3. The locations of the clumps for the best model are marked with red stars. The size of the stars is proportional to their velocity dispersion. The two big black stars show the two large dark matter clumps, their size again proportional to the velocity dispersion. *Left panel:* Configuration (A): The model consists of five freely placed galaxy sized halos in addition to DM1 and DM2. *Right panel:* Configuration (B) The model consists of ten freely placed galaxy sized halos in addition to DM1 and DM2. Both configurations place galaxy scaled clumps where the BCG and the galaxies responsible for the local splitting of S1 are located. The maps are $200'' \times 200''$ and are centered on the BCG with North being up and East being left.

Girardi et al. (1997) studied the structure of Abell 2218 using the spectroscopic data from Le Borgne et al. (1992). They found evidence for two structures (labelled MS1 and MS2) separated by 2000 km s^{-1} . The larger of these two structures, MS2, contains both the BCG and the brightest galaxy in DM2. The two structures are superimposed along the line of sight and do not correspond exactly to the clumps found for the strong lens modeling. Such superimposed structures are hard to separate using lensing as it is not sensitive to the 3D distribution of the matter.

To look for substructure associated with the BCG and #617 we have calculated the separation ($\Delta v = c(z_i - z_j)/(1 + z_i)$) between them and the other cluster members using spectroscopic data from Le Borgne et al. (1992). Out of the 50 galaxies in the catalog we found 13 with $\Delta v < 500 \text{ km s}^{-1}$ and 25 with $\Delta v < 1000 \text{ km s}^{-1}$ for the brightest galaxy in DM2 while for DM1 the numbers were 11 and 21 respectively. The $\Delta v < 500 \text{ km s}^{-1}$ cut defines two structures without any common members, while the $\Delta v < 1000 \text{ km s}^{-1}$ cut starts mixing the two groups. While all the galaxies we find with a $\Delta v < 500 \text{ km s}^{-1}$ for the BCG and #617 would belong to MS2 found by Girardi et al. (1997), the data suggests that MS2 may be further subdivided into two smaller structures, corresponding to the DM1 and DM2 found by the lens models.

The X-ray data and the distribution of the cluster members in velocity space, indicates that the bimodal mass distribution is caused by a merger. The main cluster is the one associated with the BCG (DM1 and BCG in the lensing analysis) which the second cluster associated with #617 (DM2 in the lensing analysis) has merged with, thus displacing the center of the X-ray peak from the BCG galaxy.

9. CONCLUSIONS

We have reconstructed a mass map of the rich galaxy cluster Abell 2218 using strong lensing constraints. Our

model is based on 7 multiply imaged systems and 1 arc with spectroscopic redshifts, and 6 systems without spectroscopic redshifts, of which 5 are new candidate systems proposed in this work. The model is sampled and optimized in the source plane by a Bayesian Monte Carlo Markov Chain implemented in the publicly available software Lenstool. Our best model has $\text{rms}_s = 0''.12$ in the source plane, corresponding to $\text{rms}_i = 1''.49$ in the image plane.

We have found, in agreement with previous models of Abell 2218, that the mass distribution is bimodal. We find that DM2 is larger and with a flatter core than previous models have found. The flatness of the profile near DM2 is further supported by our 'blind tests', which do not place a galaxy sized component near its center. The BCG and DM1 are the dominant component of the mass model in the inner regions ($< 100''$) of the cluster. We have analyzed the distribution of galaxies in velocity space, finding evidence for two substructures, separated by $\sim 1000 \text{ km s}^{-1}$, corresponding to DM1 and DM2. Although both the light and the X-ray contours are consistent with the mass map, the center of the X-ray emission is offset from the central peak of the BCG. We find that the X-ray data and the distribution in velocity space support the interpretation that the bimodal mass distribution arises from a cluster merger.

We have explored the degeneracy of the mass model, both those inherent to the dPIE profile and those arising from the mass model components themselves. For the dPIE, our results are in agreement with those of Jullo et al. (2007). We find that the large scale dark matter clumps are a necessary component of the model, i.e., using only the dark matter halos associated with the galaxies does not give a good fit to the data ($\text{rms}_s = 1''.62$ vs. $\text{rms}_s = 0''.12$), even if they are allowed to become much more massive.

At around $100''$ the two large scale halos contribute $\sim 85\%$ of the enclosed projected mass, while the BCG contributes $\sim 9\%$ and the remaining cluster galaxies $\sim 6\%$. We have performed 'blind tests' to check where the model requires a galaxy scale component to reproduce the lensing constraints, and find that both the BCG

and galaxies which locally perturb given multiply imaged systems are reliably reclaimed. However, we find that the inclusion of the cluster galaxies (excluding the BCG) only weakly affects the model unless they locally perturb a multiply imaged system. Assuming that mass scales with light, this shows that strong lensing constraints can reliably detect substructure, dark or luminous, if the substructure is massive or locally perturbs a system.

The accurate mass map we have presented is made available to the community and can be used to exploit Abell 2218 as a gravitational telescope, probing the high redshift universe. In this work we have fixed the cosmology, but with the increased number of constraints it may be possible to simultaneously fit it with the mass distribution. However, the bimodal structure of Abell 2218, and the remaining uncertainty in the mass model (around $\sim 1''$ in the image plane) may make it challenging to reach competitive accuracy in the derived cosmological parameters.

We thank Sébastien Bardeau and Geneviève Soucail for

giving us access to the weak lensing data. Á.E. thanks Bo Milvang-Jensen, Cécile Faure, Chloé Féron, John McKean, Kim Nilsson and Paul Vreeswijk for helpful discussions relating to this work. The Dark Cosmology Centre is funded by the Danish National Research Foundation. This work was supported by the European Community's Sixth Framework Marie Curie Research Training Network Programme, Contract No. MRTN-CT-2004-505183 "ANGLES". We thank the Danish Centre for Scientific Computing for granting the computer resources used. This research was supported in part by the National Science Foundation under Grant No. PHY99-07949. JR is grateful to the California Institute of Technology for its support. JPK acknowledges support from CNRS. KP acknowledges support from IDA. The authors recognize and acknowledge the very significant cultural role and reverence that the summit of Mauna Kea has always had within the indigenous Hawaiian community. We are most fortunate to have the opportunity to conduct observations from this mountain.

APPENDIX

A. THE DUAL PSEUDO ISOTHERMAL ELLIPTICAL MASS DISTRIBUTION

The Pseudo Isothermal Elliptical Mass Distribution (PIEMD) was first introduced by Kassiola & Kovner (1993a) and it and its variants are frequently used in lensing analysis (see e.g., Kneib et al. 1996; Keeton & Kochanek 1998; Smith et al. 2005; Limousin et al. 2007b; Natarajan et al. 2007; Richard et al. 2007). The main advantage of using the PIEMD for lensing analysis is that its potential, as well as its first and second partial derivatives, can be expressed analytically to derive deflections, distortions, amplifications and time delays for any ellipticity (see also Keeton & Kochanek (1998)). One of the most commonly used variants, and the one used in this work, is a two component PIEMD with both a core radius and a scale radius. This profile is sometimes referred to as the "truncated PIEMD" or simply the "PIEMD", although it differs in significant ways from the original PIEMD proposed by Kassiola & Kovner (1993a). An equivalent family of mass models was introduced independently by Merritt & Valluri (1996) who used it as a starting point in an investigation of the orbital properties of triaxial stellar systems. In this appendix we give a self-contained description of this profile, which we will call the 'dual Pseudo Isothermal Elliptical mass distribution' (dPIE) and present its derived quantities most relevant to lensing.

A.1. The 3D density profile

The 3D density distribution of the dPIE is

$$\rho(r) = \frac{\rho_0}{(1 + r^2/a^2)(1 + r^2/s^2)}; \quad s > a. \quad (\text{A1})$$

This distribution represents a spherical system with scale radius s , core radius a and central density ρ_0 ⁹. In the center $\rho \approx \rho_0/(1 + r^2/a^2)$ which describes a core with central density ρ_0 . The core is not strictly isothermal (in which case $\rho \sim (1 + r^2/a^2)^{-3/2}$ (Binney & Tremaine 1987)) but in the transition region, $a \lesssim r \lesssim s$, we have $\rho \sim r^{-2}$ as for the isothermal sphere. In the outer parts the density falls off as $\rho \sim r^{-4}$.

The density distribution can be rewritten as

$$\rho(r) = \frac{\rho_0}{(1 + (\eta\xi)^2)(1 + \xi^2)}; \quad \eta > 1 \quad (\text{A2})$$

with $\xi \equiv r/s$ and $\eta \equiv s/a$. Thus s acts as a scale parameter and η is a shape parameter.

A.2. The 2D density profile

The projected density, which is the relevant quantity for lensing, is given by

$$\Sigma(R) = 2 \int_R^\infty \frac{\rho(r)r}{\sqrt{r^2 - R^2}} dr = \Sigma_0 \frac{as}{s - a} \left(\frac{1}{\sqrt{a^2 + R^2}} - \frac{1}{\sqrt{s^2 + R^2}} \right) \quad (\text{A3})$$

with

$$\Sigma_0 = \pi \rho_0 \frac{as}{s + a}. \quad (\text{A4})$$

and R is the 2D radius. For a vanishing core radius it reduces to

$$\Sigma(R) = \Sigma_0 \frac{a}{R} \left(1 - \frac{1}{\sqrt{1 + s^2/R^2}} \right); \quad a \ll R \quad (\text{A5})$$

which is the surface mass profile proposed by Brainerd et al. (1996) for modeling galaxy-galaxy lensing. In this special case, a becomes a simple amplitude scaling parameter and, as shown below, s is the half-mass radius, r_h .

Defining the critical surface mass density

$$\Sigma_{\text{crit}} \equiv \frac{c^2}{4\pi G} \frac{D_S}{D_L D_{LS}}. \quad (\text{A6})$$

where D_L , D_S , and D_{LS} are the angular diameter distances to the lens, the source, and between the lens and source respectively, the convergence is

$$\kappa(R) \equiv \frac{\Sigma(R)}{\Sigma_{\text{crit}}} \quad (\text{A7})$$

and the corresponding shear is

$$\gamma(R) = \frac{\Sigma_0}{\Sigma_{\text{crit}}} \frac{as}{s - a} \left[2 \left(\frac{1}{a + \sqrt{a^2 + R^2}} - \frac{1}{s + \sqrt{s^2 + R^2}} \right) + \left(\frac{1}{\sqrt{a^2 + R^2}} - \frac{1}{\sqrt{s^2 + R^2}} \right) \right]. \quad (\text{A8})$$

⁹ We note that s corresponds to r_{cut} and a to r_{core} in previous publications (e.g. Limousin et al. (2007b); Jullo et al. (2007)), but we adopt this new notation to avoid confusion, as s is not a 'cut-off' nor a truncation radius, but the a scale radius, with $\rho \sim r^{-4}$ for $r \gg s$ (see also Section A.3 for how the mass depends on s).

A.3. *Mass relations*

The mass inside physical radius r is

$$M_{3D}(r) = 4\pi \int_0^r \rho(\tilde{r}) \tilde{r}^2 d\tilde{r} = 4\pi \rho_0 \frac{a^2 s^2}{s^2 - a^2} \left(s \arctan\left(\frac{r}{s}\right) - a \arctan\left(\frac{r}{a}\right) \right) \quad (\text{A9})$$

and the mass inside projected radius R is

$$M_{2D}(R) = 2\pi \int_0^R \Sigma(\tilde{R}) \tilde{R} d\tilde{R} = 2\pi \Sigma_0 \frac{as}{s-a} \left(\sqrt{a^2 + R^2} - a - \sqrt{s^2 + R^2} + s \right). \quad (\text{A10})$$

The dPIE has a finite total mass, given by

$$M_{\text{TOT}} = 2\pi^2 \rho_0 \frac{a^2 s^2}{a+s} = 2\pi \Sigma_0 as. \quad (\text{A11})$$

The half-mass radius defined by $M_{3D}(r_h) = M_{\text{TOT}}/2$ obeys the relation

$$s \arctan\left(\frac{r_h}{s}\right) - a \arctan\left(\frac{r_h}{a}\right) = \frac{\pi}{4}(s-a). \quad (\text{A12})$$

For $a \ll r_h, s$

$$r_h \approx s + \frac{\pi}{2}a \quad (\text{A13})$$

i.e., $r_h \geq s$ with equality for $a \rightarrow 0$. Similarly, the effective radius defined by $M_{2D}(R_e) = M_{\text{TOT}}/2$ is

$$R_e = \frac{3}{4} \sqrt{a^2 + \frac{10}{3}as + s^2} \quad (\text{A14})$$

which for $\eta^{-1} \ll 1$ ($a \ll s$) reduces to

$$R_e \approx \frac{3}{4}s + \frac{5}{4}a, \quad (\text{A15})$$

i.e., $R_e \approx (3/4)r_h$ for $a \rightarrow 0$.

A.4. *The potential*

The gravitational potential $\Psi = -\Phi$ is

$$\Psi(r) = G \int_r^\infty \frac{M_{3D}(r)}{r^2} dr = 4\pi G \rho_0 \frac{a^2 s^2}{s^2 - a^2} \left(\frac{s}{r} \arctan\left(\frac{r}{s}\right) - \frac{a}{r} \arctan\left(\frac{r}{a}\right) + \frac{1}{2} \ln\left(\frac{s^2 + r^2}{a^2 + r^2}\right) \right) \quad (\text{A16})$$

and the central potential is

$$\Psi(0) = 4\pi G \rho_0 \frac{a^2 s^2}{s^2 - a^2} \ln \frac{s}{a} \quad (\text{A17})$$

which is finite for finite a .

The projected potential is given by

$$\begin{aligned} \Psi(R) &= 2G \int_R^{R_L} \frac{M_{2D}(\tilde{R})}{\tilde{R}} d\tilde{R} \\ &= 4\pi G \Sigma_0 \frac{as}{s-a} \left(\sqrt{s^2 + R^2} - \sqrt{a^2 + R^2} + a \ln(a + \sqrt{a^2 + R^2}) - s \ln(s + \sqrt{s^2 + R^2}) \right) \\ &\quad + \text{constant}, \end{aligned} \quad (\text{A18})$$

where R_L is a limiting radius leading to the constant term. The deflection then is

$$\alpha(R) = -\frac{2}{c^2} \frac{D_{LS}}{D_S} \frac{d\Psi}{dR} = \frac{8\pi G}{c^2} \frac{D_{LS}}{D_S} \Sigma_0 \frac{as}{s-a} f(R/a, R/s) \quad (\text{A19})$$

where

$$f(R/a, R/s) \equiv \left(\frac{R/a}{1 + \sqrt{1^2 + (R/a)^2}} - \frac{R/s}{1 + \sqrt{1^2 + (R/s)^2}} \right). \quad (\text{A20})$$

A.5. Ellipticity

An elliptical projected mass distribution with ellipticity $\epsilon \equiv (A - B)/(A + B)$, where A and B are the semi major and minor axes respectively, can be introduced by substituting $R \rightarrow \tilde{R}$, with

$$\tilde{R}^2 = \frac{X^2}{(1 + \epsilon)^2} + \frac{Y^2}{(1 - \epsilon)^2}, \quad (\text{A21})$$

where X and Y are the spatial coordinates along the major and minor projected axes, respectively. With this definition of the ellipticity, the quantity $e \equiv 1 - B/A$ is related to ϵ through $e = 2\epsilon/(1 + \epsilon)$. For further details we refer to Kassiola & Kovner (1993a), and note that in their notation the model discussed in this paper corresponds to

$$J_{as,3/2}^* = \frac{as}{s - a}(a^{-1}I_{a,1/2}^* - s^{-1}I_{s,1/2}^*). \quad (\text{A22})$$

and the analytical potential is given by solving

$$\frac{\delta^2 \Phi}{\delta X^2} = \Re \frac{\delta J_{as,3/2}^*}{\delta X}; \quad \frac{\delta^2 \Phi}{\delta Y^2} = \Im \frac{\delta J_{as,3/2}^*}{\delta Y}; \quad \frac{\delta^2 \Phi}{\delta X \delta Y} = \Im \frac{\delta J_{as,3/2}^*}{\delta X} = \Re \frac{\delta J_{as,3/2}^*}{\delta Y}.$$

A.6. The dPIE in Lenstool

The dPIE is incorporated into the Lenstool software using the full elliptical expressions. The profile is given by specifying 8 parameters, its redshift z , its central position (R.A., Dec.), its ellipticity and orientation $(\hat{e}, \theta_{\hat{e}})$, the core radius a , the scale radius s and a fiducial velocity dispersion σ_{dPIE} . This fiducial velocity dispersion is related to the deflection, α , by

$$\alpha = \frac{a + s}{s} E_0 f(R/a, R/s) \quad (\text{A23})$$

where

$$E_0 = 6\pi \frac{D_{LS}}{D_S} \frac{\sigma_{\text{dPIE}}^2}{c^2}. \quad (\text{A24})$$

In terms of Σ_0 , a and s (or ρ_0 , a and s) σ_{dPIE} is given by

$$\sigma_{\text{dPIE}}^2 = \frac{4}{3} G \Sigma_0 \frac{as^2}{s^2 - a^2} = \frac{4}{3} G \pi \rho_0 \frac{a^2 s^3}{(s - a)(s + a)^2}. \quad (\text{A25})$$

The ellipticity is defined by

$$\hat{e} = \frac{A^2 - B^2}{A^2 + B^2} \quad (\text{A26})$$

corresponding to

$$e = 1 - \frac{B}{A} = 1 - \frac{\sqrt{1 - \hat{e}}}{\sqrt{1 + \hat{e}}}. \quad (\text{A27})$$

Its direction, $\theta_{\hat{e}}$, is measured anti-clockwise from the West and is related to the position angle (P.A.) by $\text{P.A.} = \theta_{\hat{e}} - 90^\circ$.

A.7. Velocity dispersions

Although σ_{dPIE} is a fiducial velocity dispersion, we wish to relate it to the measured velocity dispersion of galaxies, assuming that their profile is described by a dPIE. In this derivation we will look at the most simple case, assuming no anisotropy and a spherically symmetric dPIE profile. Even under these assumptions, the central velocity dispersion and the velocity dispersion profiles for the dPIE cannot be expressed analytically and must be computed numerically. The velocity dispersion (Binney & Tremaine 1987) is

$$\sigma^2(r) = \frac{G}{\rho(r)} \int_r^\infty \frac{M_{3D}(r)\rho(r)}{r^2} dr \quad (\text{A28})$$

and the projected (line-of-sight) velocity dispersion is

$$\sigma_P^2(R) = \frac{2G}{\Sigma(R)} \int_R^\infty \frac{M_{3D}(r)\rho(r)}{r^2} \sqrt{r^2 - R^2} dr. \quad (\text{A29})$$

We can also compute the average line-of-sight velocity dispersion inside a given radius R as

$$\langle \sigma_P^2 \rangle (R) = \frac{2\pi \int_0^R \sigma_P^2(R) \Sigma(R) R dR}{M_{2D}(R)}. \quad (\text{A30})$$

This is the quantity measured by velocity dispersion measurements, and for galaxies the relevant radius corresponds roughly to the slit width used for the spectrograph. For the galaxies in our model (see Table 2) and a spectrograph resolution of $\sim 1''$, this corresponds to an observable region of $0.01 \lesssim (R/R_e) \lesssim 2$. In Figure A1 we plot the average




FIG. 20.— The squared measured velocity dispersion ($\langle \sigma_P^2 \rangle$) vs. the squared fiducial σ_{dPIE}^2 . The ratio varies depending on both the observed radius R , and the ratio of a and s . For values similar to those found for the galaxies in the model (Table 2) we find $\sqrt{\langle \sigma_P^2 \rangle} \approx (0.85 \pm 0.10)\sigma_{\text{dPIE}}$.

line-of-sight velocity dispersion over the fiducial velocity dispersion σ_{dPIE} for different values of η . We find that for a/s values similar to those found in the modelling, $\sqrt{\langle \sigma_P^2 \rangle}/\sigma_{\text{dPIE}} \approx 0.85 \pm 0.10$ in the observable region.

Finally, to facilitate comparison with velocity dispersion profiles from different profiles, we define a 'standard' velocity dispersion σ_e . This velocity dispersion is motivated by the velocity dispersion of the isothermal profile (for which $\sigma_{\text{iso}}^2 = 2GM_{2D}(R)/R$ for all R), calculated at the effective radius R_e :

$$\sigma_e^2 \equiv 2G \langle \Sigma \rangle_e R_e \quad (\text{A31})$$

where $\langle \Sigma \rangle_e$ is the mean surface density at R_e . For the dPIE profile, we find that:

$$\frac{\sigma_e^2}{\sigma_{\text{dPIE}}^2} = \frac{2(s^2 - a^2)}{s\sqrt{a^2 + (10/3)as + s^2}} \quad (\text{A32})$$

which goes to 2 when the core radius a goes to zero. For a/s values similar to those found in the modelling $\sigma_e/\sigma_{\text{dPIE}} \approx 1.3 \pm 0.2$.

REFERENCES

- Abdelsalam, H. M., Saha, P., & Williams, L. L. R. 1998, *AJ*, 116, 1541
- Abell, G. O. 1958, *ApJS*, 3, 211
- Abell, G. O., Corwin, Jr., H. G., & Olowin, R. P. 1989, *ApJS*, 70, 1
- Bardeau, S., Soucail, G., Kneib, J.-P., Czoske, O., Ebeling, H., Hudelot, P., Smail, I., & Smith, G. P. 2007, *A&A*, 470, 449
- Binney, J. & Tremaine, S. 1987, *Galactic dynamics* (Princeton, NJ, Princeton University Press, 1987, 747 p.)
- Bolzonella, M., Miralles, J.-M., & Pelló, R. 2000, *A&A*, 363, 476
- Bradač, M., Clowe, D., Gonzalez, A. H., Marshall, P., Forman, W., Jones, C., Markevitch, M., Randall, S., Schrabback, T., & Zaritsky, D. 2006, *ApJ*, 652, 937
- Brainerd, T. G., Blandford, R. D., & Smail, I. 1996, *ApJ*, 466, 623
- Cavaliere, A. & Fusco-Femiano, R. 1978, *A&A*, 70, 677
- de Propriis, R., Stanford, S. A., Eisenhardt, P. R., Dickinson, M., & Elston, R. 1999, *AJ*, 118, 719
- Ebbels, T., Ellis, R., Kneib, J.-P., Leborgne, J.-F., Pello, R., Smail, I., & Sanahuja, B. 1998, *MNRAS*, 295, 75
- Ebbels, T. M. D., Le Borgne, J.-F., Pello, R., Ellis, R. S., Kneib, J.-P., Smail, I., & Sanahuja, B. 1996, *MNRAS*, 281, L75
- Egami, E., Kneib, J.-P., Rieke, G. H., Ellis, R. S., Richard, J., Rigby, J., Papovich, C., Stark, D., Santos, M. R., Huang, J.-S., Dole, H., Le Floc'h, E., & Pérez-González, P. G. 2005, *ApJ*, 618, L5
- Ellis, R., Santos, M. R., Kneib, J.-P., & Kuijken, K. 2001, *ApJ*, 560, L119
- Gavazzi, R. & Soucail, G. 2007, *A&A*, 462, 459
- Girardi, M., Fadda, D., Escalera, E., Giuricin, G., Mardirossian, F., & Mezzetti, M. 1997, *ApJ*, 490, 56
- Halkola, A., Seitz, S., & Pannella, M. 2006, *MNRAS*, 372, 1425
- . 2007, *ApJ*, 656, 739
- Jørgensen, I., Franx, M., Hjorth, J., & van Dokkum, P. G. 1999, *MNRAS*, 308, 833
- Jullo, E., Kneib, J.-P., Limousin, M., Elíasdóttir, Á., Marshall, P., & Verdugo, T. 2007, *ArXiv e-prints*, 706
- Kassiola, A. & Kovner, I. 1993a, *ApJ*, 417, 450
- . 1993b, *ApJ*, 417, 474
- Keeton, C. R. & Kochanek, C. S. 1998, *ApJ*, 495, 157
- Kneib, J.-P. 1993, PhD thesis, Ph. D. thesis, Université Paul Sabatier, Toulouse, (1993)
- Kneib, J.-P., Ellis, R. S., Santos, M. R., & Richard, J. 2004a, *ApJ*, 607, 697
- Kneib, J.-P., Ellis, R. S., Smail, I., Couch, W. J., & Sharples, R. M. 1996, *ApJ*, 471, 643
- Kneib, J. P., Mellier, Y., Pello, R., Miralda-Escude, J., Le Borgne, J.-F., Boehringer, H., & Picat, J.-P. 1995, *A&A*, 303, 27
- Kneib, J.-P., van der Werf, P. P., Kraiberg Knudsen, K., Smail, I., Blain, A., Frayer, D., Barnard, V., & Ivison, R. 2004b, *MNRAS*, 349, 1211
- Kochanek, C. S. 1996, *ApJ*, 466, 638
- Koopmans, L. V. E., Treu, T., Fassnacht, C. D., Blandford, R. D., & Surpi, G. 2003, *ApJ*, 599, 70
- Le Borgne, J. F., Pello, R., & Sanahuja, B. 1992, *A&AS*, 95, 87
- Le Fevre, O., Crampton, D., Lilly, S. J., Hammer, F., & Tresse, L. 1995, *ApJ*, 455, 60
- Limousin, M., Kneib, J. P., Bardeau, S., Natarajan, P., Czoske, O., Smail, I., Ebeling, H., & Smith, G. P. 2007a, *A&A*, 461, 881
- Limousin, M., Richard, J., Kneib, J. ., Jullo, E., Fort, B., Soucail, G., Elíasdóttir, A., Natarajan, P., Smail, I., Ellis, R. S., Czoske, O., Hudelot, P., Bardeau, S., Ebeling, H., & Smith, G. P. 2007b, *ApJ*, 668, 643
- Limousin, M., Sommer-Larsen, J., Natarajan, P., & Milvang-Jensen, B. 2007c, *ArXiv e-prints*
- Mandelbaum, R., Tasitsiomi, A., Seljak, U., Kravtsov, A. V., & Wechsler, R. H. 2005, *MNRAS*, 362, 1451
- Meneghetti, M., Argazzi, R., Pace, F., Moscardini, L., Dolag, K., Bartelmann, M., Li, G., & Oguri, M. 2007, *A&A*, 461, 25
- Meneghetti, M., Jain, B., Bartelmann, M., & Dolag, K. 2005, *MNRAS*, 362, 1301
- Merritt, D. & Valluri, M. 1996, *ApJ*, 471, 82
- Natarajan, P., De Lucia, G., & Springel, V. 2007, *MNRAS*, 376, 180
- Natarajan, P. & Kneib, J.-P. 1997, *MNRAS*, 287, 833
- Natarajan, P., Kneib, J.-P., & Smail, I. 2002a, *ApJ*, 580, L11
- Natarajan, P., Kneib, J.-P., Smail, I., & Ellis, R. S. 1998, *ApJ*, 499, 600
- Natarajan, P., Loeb, A., Kneib, J.-P., & Smail, I. 2002b, *ApJ*, 580, L17
- Oke, J. B., Cohen, J. G., Carr, M., Cromer, J., Dingizian, A., Harris, F. H., Labrecque, S., Lucinio, R., Schaal, W., Epps, H., & Miller, J. 1995, *PASP*, 107, 375
- Richard, J., Kneib, J.-P., Jullo, E., Covone, G., Limousin, M., Ellis, R., Stark, D., Bundy, K., Czoske, O., Ebeling, H., & Soucail, G. 2007, *ApJ*, 662, 781
- Schechter, P. L., Bailyn, C. D., Barr, R., Barvainis, R., Becker, C. M., Bernstein, G. M., Blakeslee, J. P., Bus, S. J., Dressler, A., Falco, E. E., Fesen, R. A., Fischer, P., Gebhardt, K., Harmer, D., Hewitt, J. N., Hjorth, J., Hurt, T., Jaunsen, A. O., Mateo, M., Mehlert, D., Richstone, D. O., Sparke, L. S., Thorstensen, J. R., Tonry, J. L., Wegner, G., Willmarth, D. W., & Worthey, G. 1997, *ApJ*, 475, L85
- Smail, I. & Dickinson, M. 1995, *ApJ*, 455, L99+
- Smail, I., Ellis, R. S., Dressler, A., Couch, W. J., Oemler, A. J., Sharples, R. M., & Butcher, H. 1997, *ApJ*, 479, 70
- Smail, I., Hogg, D. W., Blandford, R., Cohen, J. G., Edge, A. C., & Djorgovski, S. G. 1995, *MNRAS*, 277, 1
- Smith, G. P., Kneib, J.-P., Smail, I., Mazzotta, P., Ebeling, H., & Czoske, O. 2005, *MNRAS*, 359, 417
- Soucail, G., Kneib, J.-P., & Golse, G. 2004, *A&A*, 417, L33
- Swinbank, A. M., Smith, J., Bower, R. G., Bunker, A., Smail, I., Ellis, R. S., Smith, G. P., Kneib, J.-P., Sullivan, M., & Allington-Smith, J. 2003, *ApJ*, 598, 162
- Ziegler, B. L., Bower, R. G., Smail, I., Davies, R. L., & Lee, D. 2001, *MNRAS*, 325, 1571